



High Efficacy of Therapeutic Equine Hyperimmune Antibodies Against SARS-CoV-2 Variants of Concern

Andres Moreira-Soto^{1,2*}, Mauricio Arguedas³, Hebleen Brenes⁴, Willem Buján^{5,6}, Eugenia Corrales-Aguilar², Cecilia Díaz^{3,5}, Ann Echeverri⁶, Marietta Flores-Díaz³, Aarón Gómez³, Andrés Hernández³, María Herrera³, Guillermo León³, Román Macaya⁶, Arne Kühne¹, José Arturo Molina-Mora², Javier Mora², Alfredo Sanabria⁶, Andrés Sánchez³, Laura Sánchez³, Álvaro Segura³, Eduardo Segura³, Daniela Solano³, Claudio Soto⁴, Jennifer L. Stynoski³, Mariángela Vargas³, Mauren Villalta³, Chantal B. E. M. Reusken⁷, Christian Drosten¹, José María Gutiérrez³, Alberto Alape-Girón^{3,5} and Jan Felix Drexler^{1,8}

OPEN ACCESS

Edited by:

Pragya Dhruv Yadav,
National Institute of Virology
(ICMR), India

Reviewed by:

Tomer Israely,
Israel Institute for Biological Research
(IIBR), Israel
Daniele Focosi,
Pisana University Hospital, Italy

*Correspondence:

Andres Moreira-Soto
andres.moreira-soto@charite.de

Specialty section:

This article was submitted to
Infectious Diseases - Surveillance,
Prevention and Treatment,
a section of the journal
Frontiers in Medicine

Received: 03 July 2021

Accepted: 10 August 2021

Published: 06 September 2021

Citation:

Moreira-Soto A, Arguedas M,
Brenes H, Buján W,
Corrales-Aguilar E, Díaz C,
Echeverri A, Flores-Díaz M, Gómez A,
Hernández A, Herrera M, León G,
Macaya R, Kühne A, Molina-Mora JA,
Mora J, Sanabria A, Sánchez A,
Sánchez L, Segura Á, Segura E,
Solano D, Soto C, Stynoski JL,
Vargas M, Villalta M, Reusken CBEM,
Drosten C, Gutiérrez JM,
Alape-Girón A and Drexler JF (2021)
High Efficacy of Therapeutic Equine
Hyperimmune Antibodies Against
SARS-CoV-2 Variants of Concern.
Front. Med. 8:735853.
doi: 10.3389/fmed.2021.735853

¹ Institute of Virology, Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany, ² Centro de Investigación en Enfermedades Tropicales (CIET), Facultad de Microbiología, Universidad de Costa Rica, San Jose, Costa Rica, ³ Instituto Clodomiro Picado, Facultad de Microbiología, Universidad de Costa Rica, San Jose, Costa Rica, ⁴ Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud, Ministry of Health, Tres Ríos, Costa Rica, ⁵ School of Medicine, Universidad de Costa Rica, San Jose, Costa Rica, ⁶ Caja Costarricense del Seguro Social, San Jose, Costa Rica, ⁷ Centre for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, Netherlands, ⁸ German Centre for Infection Research (DZIF), Associated Partner Charité-Universitätsmedizin Berlin, Berlin, Germany

SARS-CoV-2 variants of concern show reduced neutralization by vaccine-induced and therapeutic monoclonal antibodies; therefore, treatment alternatives are needed. We tested therapeutic equine polyclonal antibodies (pAbs) that are being assessed in clinical trials in Costa Rica against five globally circulating variants of concern: alpha, beta, epsilon, gamma and delta, using plaque reduction neutralization assays. We show that equine pAbs efficiently neutralize the variants of concern, with inhibitory concentrations in the range of 0.146–1.078 $\mu\text{g/mL}$, which correspond to extremely low concentrations when compared to pAbs doses used in clinical trials. Equine pAbs are an effective, broad coverage, low-cost and a scalable COVID-19 treatment.

Keywords: equine antibodies, SARS-CoV-2, therapy, variant of concern, PRNT titers 50, neutralization test, COVID-19

SARS-CoV-2 causes coronavirus infectious disease 19 (COVID-19), which leads to either critical illness or death in 5% of patients (1). COVID-19 prevention and treatment options include vaccines, antivirals, and antibody formulations. A wide array of vaccine platforms have shown efficacy in preventing severe disease, but universal access is limited in many resource-limited settings lacking sufficient vaccine coverage (2). Even though there are more than 300 therapeutic drugs in clinical trials, few have proven effective, such as dexamethasone (1, 3). Direct-acting antivirals like Remdesivir are most effective if given very early in the course of the disease, require supplementary oxygen therapy and are very costly at 2,000–3,000 USD per treatment, limiting universal access (4). The use of monoclonal antibodies (mAbs) are safe alternatives shown to enhance viral clearance (5), but their large-scale production is challenging and costly, at around 1,500–6,500 USD per treatment. Polyclonal antibodies (pAbs), either homologous in the case of convalescent plasma and hyperimmune sera, or heterologous such as equine hyperimmune sera, constitute a proven alternative. Convalescent plasma is readily used as COVID-19 therapy

due to its rapid capacity of deployment, decade long proven efficacy against emerging diseases such as Ebola and influenza (6), and affordability, at 350–1,000 USD per treatment. Another advantage of convalescent plasma is the use of routine blood donors or follow-up sera of discharged patients, which leads to the production of antibodies against the circulating pathogen, reducing the possibility of immune evasion (6). Nevertheless, patients with mild symptomatology may develop low-titer antibodies as observed for other emerging infectious diseases (6). To overcome this obstacle, hyperimmune globulins can be used, which are prepared from the pooling of many donors. However, both convalescent plasma and hyperimmune sera are donor-dependent, require strict donor rigorous testing for both blood-borne pathogens and high levels of neutralizing anti-SARS-CoV-2 antibodies, all of which might not be readily available in blood bank systems in many developing countries (5, 7). Another low-cost alternative are formulations of intact or fragmented equine polyclonal antibodies (pAbs), widely used for decades as therapies against some viral infections or as antivenoms (8).

We and others have previously shown that horses can be efficiently immunized with different SARS-CoV-2 antigens to yield high quantities of purified pAbs that are 50–80 times more potent than convalescent plasma for virus neutralization (9, 10). A formulation of equine polyclonal F(ab')₂ fragments against the receptor binding domain (RBD) of SARS-CoV-2 was tested in a multi-center, double-blind, placebo-controlled phase II/III clinical trial showing that it is well tolerated and leads to clinical improvement of hospitalized patients with moderate to severe COVID-19 (11). Additionally, there is an ongoing randomized, multi-center, double-blind, placebo-controlled, dose-finding, phase IIb/III clinical trial (NCT04838821) at hospitals of the Costa Rican Social Security Fund testing equine pAbs formulations to treat moderate and severe COVID-19 cases.


However, pre-clinical data of equine hyperimmune pAbs are only available for early SARS-CoV-2 isolates, whereas such data are lacking for recent and globally circulating variants, considered of concern (VoC) due to their increased transmissibility. VoC alpha, beta, epsilon, gamma and delta (<https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>) (lineage designations in Pango/Nextrain: B.1.1.7/501Y.V1 first detected in the United Kingdom, B.1.351/501Y.V2 first detected in South Africa, P.1/501Y.V3 first detected in Brazil/Japan, B.1.427/B.1.429 first detected in the US/California and B.1.617.2/S:478K first detected in India) exhibit a substantial reduction of neutralization by therapeutic mAbs or by antibodies present in the plasma of vaccinated or convalescent individuals (12, 13).

Here we report the results of a plaque reduction neutralization assay (PRNT) against VoC for our purified equine pAbs formulations. PRNT were performed as follows. Briefly, VeroE6 cells (3.25×10^5 cells/ml) were seeded in 24-well plates and incubated overnight. Equine pAbs formulations were mixed in equal parts with a virus solution containing 20 PFU/well previously titrated in the same cells. The experiment was performed in triplicate, and six wells were incubated only with the virus solution containing 20 PFU/well as positive control. The antibody–virus solution was incubated at 37°C for 1 hour

and added to the cells. After 1 hour at 37°C, supernatants were discarded, and cells were supplemented with 1.2% Avicel solution in DMEM. After 3 days at 37°C, supernatants were removed, and the 24-well plates were fixed and inactivated using a 6% formaldehyde/PBS solution and stained with crystal violet, and plaques were counted.

The two formulations contain antibodies either against the SARS-CoV-2 recombinant S1 protein (called anti-S1; produced in baculovirus insect cells), or SEM mosaic (called anti-mix; an *E. coli* derived recombinant protein containing the S, E, and M immunodominant regions) derived from the strain Wuhan-Hu-1, Accession Number: YP_009724390 (Native Antigen Company, Oxford, United Kingdom), purified using caprylic acid precipitation method (10). Both formulations effectively neutralized the five VoC and an early isolate of the virus (Germany/Gisaid_EPI_ISL_406862) at similar low inhibitory concentrations (IC₅₀ range for anti-S1 formulation: 0.206–1.078 µg/mL; and for the anti-mix formulation: 0.146–0.8359 µg/mL; **Supplementary Figure 1**; IC₅₀ dose-response curves are shown in the Technical Annex). The highest IC₅₀ corresponds to the delta VoC (**Figure 1**). Differences between potencies were statistically significant for the anti-S1 formulation (sum-of-squares F test; $p < 0.01$). This difference was only observed when the delta variant was added to the dataset (sum-of-squares F test without the delta VoC; $p = 0.9$). Previously, the delta VoC has shown resistance to therapeutic monoclonal antibodies due to binding impairment to the spike protein, and fourfold reduced sensitivity in neutralization tests performed in sera of convalescent individuals, suggesting increased immune evasion (14). Therefore, it is expected that a higher dose is needed for neutralization of the delta VoC, in agreement with our data. For the anti-mix formulation, the differences between potencies against tested VoC and early SARS-CoV-2 isolates were not statistically significant (sum-of-squares F test; $p = 0.3$). Notably, IC₅₀ values against the five tested VoC of both formulations are extremely low when compared to pAbs doses used by other groups in patients enrolled in clinical trials (4 mg/kg) (11), even at the upper estimates of the 95% confidence intervals, reaching a maximum of 13.89 µg/mL for the beta VoC (**Figure 1**). Hypothetically, hyperimmunization protocols with potent adjuvants lead to a strong immune response in the horses. Therefore, antibodies likely recognize a diverse variety of epitopes in the viral proteins, showing higher binding affinity that ensure a wider recognition and neutralization than in the case of plasma from vaccinated people or convalescent plasma.

The use of equine pAbs as potential COVID-19 therapy shows several limitations including the need for an early administration during the course of the disease, which is a limitation of all antibody-based therapies, and the risk of adverse reactions to equine immunoglobulins, including serum sickness, owing to the heterologous nature of the preparations. Nevertheless, such reactions can be readily managed pharmacologically, as evidenced by the long-standing experience with the use of equine-derived antivenoms. Our data underscore the high potential of equine pAbs for treatment of COVID-19. As more VoC emerge, further studies should



evaluate whether this wide cross-neutralization between SARS-CoV-2 variants is maintained. Shifting antivenom platforms to produce equine pAbs, laboratories in both developed and developing countries that have been manufacturing and distributing safe and standardized antivenoms for decades could rapidly fill the gaps in global demand for therapies that are both effective against VoC and affordable to low- and middle-income countries.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

AM-S, MA, HB, WB, EC-A, CDí, AE, MF-D, AG, AH, MH, GL, RM, JM-M, JM, ASana, ASánc, LS, ÁS, ES, DS, CS, JS, MVa, MVi, AA-G, JG, and JD conceived, planned and performed the experiments. AM-S and AK carried out the BSL-3 experiments. JD, AA-G, JG, and AM-S planned and carried out the analyses. AM-S, JD, CDr, CR, AA-G, and JG contributed to the interpretation of the results. AM-S, JD, AA-G, and JG took the lead in writing the manuscript. All authors provided critical feedback, helped shape the research, analysis, and manuscript.

FUNDING

This work was supported by the Global Centres for Health and Pandemic Prevention from the German academic exchange services (DAAD) (Grant agreement: 57592642), the Vicerrectoría de Investigación, Universidad de Costa Rica (Project 741-C0-198), and the Banco Centroamericano de Integración Económica (BCIE).

ACKNOWLEDGMENTS

We acknowledge support from the German Research Foundation (DFG) and the Open Access Publication Fund of Charité - Universitätsmedizin Berlin.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmed.2021.735853/full#supplementary-material>

Supplementary Figure 1 | IC₅₀ dose-response curves of the antibody preparations to SARS-CoV-2 early isolates and variants of concern named using WHO and Pango/Nextrain designations. The Y axis denotes the mean plaque forming units (PFU) per milliliter in triplicate. The X axis denotes the Log₁₀ concentration of the Anti-S1 and the Anti-Mix (combination of S1, N and SEM mosaic protein of Wuhan-Hu-1, Accession N YP_009724390.1) formulations.

REFERENCES

- Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol.* (2021) 19:141–54. doi: 10.1038/s41579-020-00459-7
- Bollyky TJ, Gostin LO, Hamburg MA. The equitable distribution of COVID-19 therapeutics and vaccines. *JAMA.* (2020) 323:2462–3. doi: 10.1001/jama.2020.6641
- The RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with COVID-19. *N Engl J Med.* (2020) 384:693–704. doi: 10.1056/NEJMoa2021436
- Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the treatment of COVID-19—final report. *N Engl J Med.* (2020) 383:1813–26. doi: 10.1056/NEJMoa2007764
- The Lancet Haematology. The resurgence of convalescent plasma therapy. *Lancet Haematol.* (2020) 7:e353. doi: 10.1016/S2352-3026(20)30117-4
- Focosi D, Anderson AO, Tang JW, Tuccori M. Convalescent plasma therapy for COVID-19: state of the art. *Clin Microbiol Rev.* (2020) 33:00072–20. doi: 10.1128/CMR.00072-20
- Focosi D, Franchini M. COVID-19 convalescent plasma therapy: hit fast, hit hard! *Vox Sang.* (2021) doi: 10.1111/vox.13091
- Gutiérrez JM, Lomonte B, Sanz L, Calvete JJ, Pla D. Immunological profile of antivenoms: preclinical analysis of the efficacy of a polyspecific antivenom through antivenomics and neutralization assays. *J Proteomics.* (2014) 105:340–50. doi: 10.1016/j.jprot.2014.02.021
- Zylberman V, Sanguineti S, Pontoriero AV, Higa SV, Cerutti ML, Morrone Seijo SM, et al. Development of a hyperimmune equine serum therapy for COVID-19 in Argentina. *Medicina.* (2020) 3:1–6.
- León G, Herrera M, Vargas M, Arguedas M, Sánchez A, Segura Á, et al. Development and characterization of two equine formulations towards SARS-CoV-2 proteins for the potential treatment of COVID-19. *Sci Rep.* (2021) 11:9825. doi: 10.1038/s41598-021-89242-z
- Lopardo G, Belloso WH, Nannini E, Colonna M, Sanguineti S, Zylberman V, et al. RBD-specific polyclonal F(ab')₂ fragments of equine antibodies in patients with moderate to severe COVID-19 disease: a randomized, multicenter, double-blind, placebo-controlled, adaptive phase 2/3 clinical trial. *EClinical Medicine.* (2021) 34:11. doi: 10.1016/j.eclinm.2021.100843
- Chen RE, Zhang X, Case JB, Winkler ES, Liu Y, Van Blargan LA, et al. Resistance of SARS-CoV-2 variants to neutralization by monoclonal and serum-derived polyclonal antibodies. *Nat Med.* (2021) 27:717–26. doi: 10.1038/s41591-021-01294-w
- Focosi D, Tuccori M, Baj A, Maggi F. SARS-CoV-2 variants: a synopsis of *in vitro* efficacy data of convalescent plasma, currently marketed vaccines, and monoclonal antibodies. *Viruses.* (2021) 13:1211. doi: 10.3390/v13071211
- Planas D, Veyer D, Baidaliuk A, Staropoli I, Guivel-Benhassine F, Rajah MM, et al. Reduced sensitivity of SARS-CoV-2 variant delta to antibody neutralization. *Nature.* (2021) 596:276–80. doi: 10.1038/s41586-021-03777-9

Conflict of Interest: Several authors of this manuscript are employees of Instituto Clodomiro Picado from University of Costa Rica, a public research institute with no commercial interests, where these antibody formulations were developed, and where eventually will be manufactured for use in the Costarican public health system.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Moreira-Soto, Arguedas, Brenes, Buján, Corrales-Aguilar, Díaz, Echeverri, Flores-Díaz, Gómez, Hernández, Herrera, León, Macaya, Kühne, Molina-Mora, Mora, Sanabria, Sánchez, Sánchez, Segura, Segura, Solano, Soto, Stynoski, Vargas, Villalta, Reusken, Drosten, Gutiérrez, Alape-Girón and Drexler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.